

MATH 5C – SAMPLE FINAL EXAM

- (1) The position vector of a particle is $\mathbf{r}(t) = e^t \mathbf{i} + t \mathbf{j}$, $t \geq 0$.
- Find equations of the tangent line at the point $(1,0)$
 - Sketch the graph of $\mathbf{r}(t)$ showing direction of increasing t .
 - Set up only, an integral to find the length of the curve from $t=0$ to $t=2$.

- (2) Evaluate the integral expression:

$$\int_0^3 \int_{y^2}^9 y \cos(x^2) dx dy$$

(Hint: You *may* want to try reversing the order of integration, although it is not necessary)

- (3) Find parametric equations of the line that passes through the point of intersection of L_1 and L_2 and is orthogonal to both L_1 and L_2 where L_1 and L_2 are the lines given by:

$$L_1 \begin{cases} x = 2t - 1 \\ y = 1 - t \\ z = 3t \end{cases} \quad L_2 \begin{cases} x = 1 + s \\ y = 2s \\ z = 3 - 2s \end{cases}$$

- (4) Find the point of intersection (if any) of the tangent lines to the curve $\mathbf{r}(t) = \langle \cos t, \cos t, \sqrt{2} \sin t \rangle$ at the points where $t = 0$ and $t = \pi/4$

- (5) SET UP BUT DO NOT EVALUATE: integrals as specified to find the volume of the wedge in the first octant cut from the cylinder $x^2 + y^2 = 4$, the xy plane, and the plane $z = y$.

a) Show sketch.

b) Triple integral - dx first.

c) Triple integral - dy first.

d) Double integral.

- (6) Given $f(x,y) = x^2 e^y$,

a) Find $D_{\mathbf{u}} f(-2,0)$ in the direction of $\vec{\mathbf{a}} = \langle 4, 3 \rangle$.

b) Find the maximum value of the directional derivative at $(-2,0)$ and the direction in which it occurs.


(7) SET UP BUT DO NOT EVALUATE: integrals as specified to find the volume enclosed by the cone $z = \sqrt{x^2 + y^2}$ and the sphere $x^2 + y^2 + z^2 = 8$.

- a) Sketch the solid
- b) Triple integral - cylindrical coordinates.
- c) Triple integral - spherical coordinates.
- d) Triple integral - rectangular coordinates .
- e) Now actually find the volume.

(8) Find the dimensions of a rectangular box, open at the top, having a volume of 108 ft^3 and requiring the least amount of material for its construction.

(9) Given the vector field $\vec{F}(x,y) = \langle y^3+1, 3xy^2+1 \rangle$ and the semicircular path C given by $\vec{r}(t) = (1 - \cos t) \vec{i} + \sin t \vec{j}$, $0 \leq t \leq \pi$ as shown,

- a) Show that \vec{F} is a conservative vector field.
- b) Find the potential function $f(x,y)$ such that $\vec{\nabla}f = \vec{F}$.
- c) Find $\int_C \vec{F} \cdot d\vec{r}$ using f .
- d) Find $\int_C \vec{F} \cdot d\vec{r}$ using a different method.



(10) Given $\vec{F}(x,y,z) = (x - y) \vec{i} + (y - z) \vec{j} + (z - x) \vec{k}$ and C is the intersection of the paraboloid $z = x^2 + y^2$ and the sphere $x^2 + y^2 + z^2 = 2$, counter clockwise when viewed from above, find $\int_C \vec{F} \cdot d\vec{r}$ two ways:

- (a) directly, and
- (b) using an appropriate theorem.

ANSWERS:

(1) (a) $\begin{cases} x = 1+t \\ y = t \end{cases}$ (b) graph of $y = \ln(x)$ from point $(1,0)$ to right (c) $\int_0^2 \sqrt{e^{2t} + 1} dt$

(2) $\frac{1}{4} \sin 81$ (3) $x=1-4t, y=7t, z=3+5t$ (4) $(1, 1, 2 - \sqrt{2})$

(5) b) $\int_0^2 \int_0^y \int_0^{\sqrt{4-y^2}} dx dz dy$

c) $\int_0^2 \int_0^{\sqrt{4-x^2}} \int_z^{\sqrt{4-x^2}} dy dz dx$

d) $\int_0^2 \int_0^{\sqrt{4-x^2}} y dy dx$

(6) (a) $-4/5$ (b) In direction of gradient $\langle -4, 4 \rangle$, value $4\sqrt{2}$.

(7) b) $\int_0^{2\pi} \int_0^2 \int_r^{\sqrt{8-r^2}} r dz dr d\theta$

c) $\int_0^{2\pi} \int_0^{\pi/4} \int_0^{\sqrt{8}} \rho^2 \sin \phi d\rho d\phi d\theta$

d) $4 \int_0^2 \int_0^{\sqrt{4-x^2}} \int_{\sqrt{x^2+y^2}}^{\sqrt{8-x^2-y^2}} dz dy dx$

e) $\frac{32\pi}{3}(\sqrt{2}-1)$

(8) Minimize $2yz+2xz+xy$ subject to $xyz=108$. Ans: 6ft by 6 ft by 3 ft high.

(9) (b) $f(x,y) = xy^3+x+y+C$ (c) 2 (d) 2

(10) π